

LAW OFFICES
MILLEN, WHITE, ZELANO & BRANIGAN, P.C.
ARLINGTON COURTHOUSE PLAZA I
SUITE 1100
2200 CLARENDON BOULEVARD
ARLINGTON, VIRGINIA 22201

TELEPHONE (703) 243-6333
CABLE USALAW
INTL TELEX 64191
TELECOPIER (703) 243-6410

Atty's Docket No. EPROV-15

JCS55 U.S. PTO
09/551405
04/17/00

Applicant(s) : Hans Rudolf MÜLLER ET AL.

For : STABLE CRYSTALLINE SALTS OF 5-METHYLTETRAHYDROFOLIC ACID

THE COMMISSIONER OF PATENTS & TRADEMARKS
Washington, D.C. 20231

SUBMISSION OF APPLICATION UNDER 37 CFR 1.53

Sir:

Herewith is the above-identified application for Letters Patent including:

Applicant(s) Name(s): 1) Hans Rudolf MÜLLER, Schaffhausen, Switzerland
2) Rudolf MOSER, Schaffhausen, Switzerland
3) Thomas EGGER, Effretikon, Switzerland

Assignee (by unrecorded assignment): EPROVA AG

Pages of Application: Specification - 11
Claims - 2
Abstract - 1
Sheets of Drawings - 5
Tables 4

NO DECLARATION IS ATTACHED

☒ Preliminary Amendment

☐ Information Disclosure Statement

☒ The benefit under 35 USC 119 is claimed of the filing date of:

SWISS APPLICATION NO. 695/99, filed April 15, 1999

☒ A certified copy of the priority document(s) is attached.

Respectfully submitted,

MILLEN, WHITE, ZELANO & BRANIGAN, P.C.

By: I. William Millen

I. William Millen (19,544)
Attorney for Applicants

IWM:djc

Date: April 17, 2000

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of : BOX PATENT APPLICATIONS

Hans Rudolf MÜLLER et al. : Examiner: Unassigned

Serial No.: Unassigned : Group Art Unit: Unassigned

Filed: April 17, 2000 :

For: STABLE CRYSTALLINE SALTS OF 5-METHYLTETRAHYDROFOLIC ACID

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Prior to examination, Applicants wish to amend the above-identified application as indicated below.

IN THE SPECIFICATION

Please amend the specification as follows:

Page 1, before the first paragraph, insert

--Field of the Invention--.

Page 1, before line 8, insert

--Background of the Invention--.

Page 2, before line 4, insert

--Summary of the Invention--.

Page 2, Line 9: Delete "limitations" and insert --limitation--.

Line 21: Delete "Figure 1 to Figure 4, and".

Line 31: Delete "Salts" and insert --Other salts--.

Page 2, bottom of page, insert

--Brief Description of the Drawings

Figures 1-4 are self-explanatory X-ray powder diffraction diagrams of the 4 crystalline modifications of the calcium salts of the invention, and Figure 5 is a diagram of the amorphous salt.--

IN THE CLAIMS

Please amend the claims as follows:

1. (Amended) [Crystalline salts] A crystalline salt of 5-methyl-(6R,S)-, -(6S)- [and] or -(6R)-tetrahydrofolic acid said crystalline salt having a water of crystallization of at least one equivalent per equivalent of 5-methyltetrahydrofolic acid.

2. (Amended) [Crystalline salts] A crystalline salt according to claim 1, of 5-methyl-(6S)- [and] or -(6R)-tetrahydrofolic acid.

3. (Amended) [The] A crystalline calcium salt according to claim 1, of 5-methyl-(6S)- and -(6R)-tetrahydrofolic acid having ≥ 3 equivalents of water.

4. (Amended) [The] A crystalline calcium salt according to claim 1, of 5-methyl-(6S)-tetrahydrofolic acid with 2 theta values of 6.5, 13.3, 16.8 and 20.1 (Type I).

5. (Amended) [The] A crystalline calcium salt according to claim 1, of 5-methyl-(6S)-tetrahydrofolic acid with 2 theta values of 5.3, 6.9, 18.7 and 21.1 (Type II).

6. (Amended) [The] A crystalline calcium salt according to claim 1, of 5-methyl-(6S)-tetrahydrofolic acid with 2 theta values of 6.8, 10.2, 15.4 and 22.5 (Type III).

7. (Amended) [The] A crystalline calcium salt according to claim 1, of 5-methyl-(6S)-tetrahydrofolic acid with 2 theta values of 6.6, 15.9, 20.2 and 22.5 (Type IV).

8. (Amended) A method of producing crystalline salts of 5-methyl-(6R,S)-, - (6S)- and 5-methyl-(6R)-tetrahydrofolic acid, [characterised in that salts] comprising subjecting a salt of 5-methyl-(6R,S)-, -(6S)- or -(6R)-tetrahydrofolic acid [are crystallised from] in a polar medium [after] to a thermal treatment, at a temperature above 60°C, and thereafter crystallizing said salt from the resultant heated solution.

Please cancel claim 9 without prejudice or disclaimer.

Claim 10, Line 1: Delete "characterised in that" and insert --wherein the--.

Claim 11, Line 1: Delete "characterised in that" and insert --wherein the--.

Claim 12, Line 1: Delete "characterised in that" and insert --wherein the--.

Claim 13, Line 1: Delete "or 12, characterised in that" and insert --wherein the--.

Delete claims 14 and 15 without prejudice or disclaimer.

Please add the following new claims:

-- **16.** A method according to claim 8, wherein said salt is an alkaline earth salt.

17. A method according to claim 8, wherein said salt is calcium.

18. A method of producing the crystalline salt of claim 5, comprising drying sufficiently a crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid with 2 theta values of 6.5, 13.3, 16.8 and 20.1 (Type I).

19. A method of producing the crystalline salt of claim 6, comprising subjecting to sufficient thermal treatment at above 90°C, a crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid with 2 theta values of 6.5, 13.3, 16.8 and 20.1 (Type I).

20. A method of producing the crystalline salt of claim 7, comprising subjecting to sufficient thermal treatment at above 95°C, a crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid with 2 theta values of 6.5, 13.3, 16.8 and 20.1 (Type I).--

REMARKS

A principle purpose of this Preliminary Amendment is to eliminate multiply dependent claims and the fee associated therewith, Applicants reserving the right to reintroduce claims to canceled combined subject matter. Also, where 2 theta values are not set forth, the water of crystallization is added to the claims.

Respectfully submitted,



I. William Millen (Reg. No. 19,544)
Attorney for Applicant(s)

MILLEN, WHITE, ZELANO & BRANIGAN, P.C.
Arlington Courthouse Plaza I
2200 Clarendon Boulevard, Suite 1400
Arlington, VA 22201
(703) 812-5325
Internet Address: millen@mwzb.com

Filed: April 17, 2000

I:\WM(pdr)\K:\PAT\eprov\15\PA.wpd

Stable crystalline salts of 5-methyltetrahydrofolic acid

This invention relates to crystalline salts of N-[4-[(2-amino-1,4,5,6,7,8hexahydro-4-oxo-5-methyl-(6S)-, -(6R)- and -(6R,S)-pteridiny]methyl]amino]benzoyl-L-glutamic acid (hereinafter called salts of 5-methyltetrahydrofolic acid), to the use thereof, and to a method of producing them.

Tetrahydrofolates are predominantly used as 5-formyltetrahydrofolic acid and the salts thereof (leucovorin) or as 5-methyltetrahydrofolic acid and the salts thereof, for the treatment of megaloblastic folic acid anaemia, as an antidote for increasing the compatibility of folic acid antagonists, particularly of aminopterin and methotrexate in cancer therapy ("antifolate rescue"), for increasing the therapeutic effect of fluorinated pyrimidines and for the treatment of autoimmune diseases such as psoriasis and rheumatoid arthritis, for increasing the compatibility of certain antiparasitic formulations, for increasing the compatibility of certain antiparasitic formulations, for instance trimethoprim-sulfamethoxazole, and for reducing the toxicity of dideazetetrahydrofolates in chemotherapy. 5-methyltetrahydrofolic acid is used in particular as a drug and as a food additive, as a vitamin preparation, for the prevention of neural tube defects, for the treatment of depressive illnesses, and for influencing the homocysteine level.

5-methyltetrahydrofolic acid and salts thereof are extremely unstable, and in particular are highly susceptible to oxidation [see also A.L. Fitzhugh, Pteridines 4 (4), 187-191 (1993) in this respect] and are therefore difficult to produce at a level of purity which is acceptable for a pharmaceutical active ingredient or a food additive.

Various methods, such as excluding oxygen as completely as possible or the addition of antioxidants such as ascorbic acid or reduced L-glutathione, have been employed in order to overcome the instability of 5-methyltetrahydrofolic acid. However, it is scarcely possible completely to exclude oxygen during use, and even then this is only possible at very considerable cost, and the addition of antioxidants is likewise not always possible. Accordingly, it has not been possible hitherto to identify a commer-

cially feasible method which is suitable for the production of salts of 5-methyltetrahydrofolic acid which are satisfactorily stable and which are of high purity.

Surprisingly, it has now been found that salts of 5-methyltetrahydrofolic acid which exhibit high chemical purity and excellent stability can be obtained by crystallising the corresponding salt from a polar medium after subjecting the solution to thermal treatment at a temperature above 60°C. The highly crystalline salts of 5-methyltetrahydrofolic acid which are thus obtained are stable at room temperature, practically without limitations. They are suitable as a constituent or as a starting material for the production of drug forms or food additives.

Accordingly, the present invention relates to crystalline salts of 5-methyltetrahydrofolic acid. Alkaline earth salts, particularly the calcium salt, are preferably used as the salts of 5-methyltetrahydrofolic acid for crystallisation. These crystalline salts of 5-methyltetrahydrofolic acid exhibit a purity, which has never been achieved hitherto, of > 98%, together with a stability, with respect to the initial value thereof and which has never been achieved hitherto, of > 98% after storage for 6 months in air at 25°C and 60% relative atmospheric humidity. The crystalline calcium salts of 5-methyl-(6S)-tetrahydrofolic acid exist in four different crystalline modifications (Type I, Type II, Type III and Type IV) and exhibit sharp bands when subjected to X-ray powder diffraction measurements (see Figure 1 to Figure 4, and Table 1 to Table 4 in this respect). Selected 2 theta values for the different crystalline modifications are 6.5, 13.3, 16.8 and 20.1 (Type I); 5.3, 6.9, 18.7 and 21.1 (Type II); 6.8, 10.2, 15.4 and 22.5 (Type III); and 6.6, 15.9, 20.2 and 22.5 (Type IV). Crystalline calcium salts of 5-methyltetrahydrofolic acid have a content of water of crystallisation of at least 1 equivalent of water per 1 equivalent of 5-methyltetrahydrofolic acid. Thus the Type I modification typically contains ≥ 3 equivalents of water, the Type II modification typically contains ≤ 2 equivalents of water and the Type III and Type IV modifications typically contain ≤ 5 equivalents of water.

Salts of 5-methyl-(6R)-tetrahydrofolic acid and salts of 5-methyl-(6R,S)-tetrahydrofolic acid can likewise be obtained in highly crystalline form.

The present invention further relates to a method of producing highly crystalline salts of 5-methyltetrahydrofolic acid, which is characterised in that the corresponding salt of 5-methyltetrahydrofolic acid is crystallised. In this method, crystallisation of salts of 5-methyltetrahydrofolic acid is effected from a polar medium after thermal treatment at a temperature above 60°C, particularly above 85°C.

Substances which are particularly suitable as the polar medium include water or a mixture of water and an organic solvent which is miscible with water, such as water-soluble alcohols, e.g. methanol, ethanol, n-propanol, iso-propanol or ethylene glycol, a low molecular weight aliphatic water-soluble carboxylic acid e.g. formic acid, acetic acid or lactic acid, or water-soluble amides e.g. formamide, dimethylformamide, dimethylacetamide, 1-methylpyrrolidone, 2-methylpyrrolidone or 2-piperidinone. There are no particular restrictions with regard to the type of solvent used and with regard to the mixture ratio, since crystalline salts of 5-methyltetrahydrofolic acid generally exhibit solubilities which are lower than those of the corresponding amorphous forms.

Crystallisation is preferably effected from solutions. It is also possible to effect crystallisation from a suspension, however.

Aside from calcium salts, still further salts of 5-methyl-(6R,S) or (6R) or (6S) hydrofolic acids include but are not limited to other alkaline earth salts, for example, magnesium can be obtained in highly crystalline forms.

The different crystalline modifications can be converted into one another by further thermal treatments at temperatures above 60°C. Thus Type I, which is produced by crystallisation from a polar medium after thermal treatment at a temperature above 60°C, can be converted into Type II by drying sufficiently, e.g. under vacuum at 70°C, can be converted into Type III by sufficient thermal treatment at a temperature above 90°C, and can be converted into Type IV by sufficient thermal treatment at a temperature above 95°C. Type II can be converted into Type I again by adding water to the crystals, e.g. by treatment with water in a humidity cabinet at 90°C.

Crystallisation of the salts of 5-methyltetrahydrofolic acid occurs spontaneously or is effected by seeding with the corresponding crystalline salt of 5-methyltetrahydrofolic acid.

A suitable, preferred starting material for crystallisation is pure, amorphous or crystalline 5-methyl-(6S)- or -(6R)-tetrahydrofolic acid. Racemic 5-methyl-(6R,S)-tetrahydrofolic acid can also be used, however, as can enriched 5-methyl-(6S)-, -(6R)- or -(6R,S)-tetrahydrofolic acid.

5

By using amorphous or partly crystalline, optically pure 5-methyltetrahydrofolic acid or salts thereof as the starting material for crystallisation, essentially crystalline salts of 5-methyltetrahydrofolic acid of a purity which has never been achieved hitherto, together with a stability which has never been achieved hitherto, are obtained by the method described here.

10

The present invention also relates to the use of highly crystalline salts of 5-methyltetrahydrofolic acid as a constituent for the production of drugs or food additive substances or for the production of other tetrahydrofolic acid derivatives, since, on account of their excellent stability in solid form, crystalline salts of 5-methyltetrahydrofolic acid are of a very good quality which remains constant with time, practically without limits. The present invention also relates to preparations containing highly crystalline salts of 5-methyltetrahydrofolic acid. These preparations are produced by known methods. They are employed analogously to the use of known substances from the field of tetrahydrofolates, such as 5-formyltetrahydrofolic acid (leucovorin) for example.

15

20

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

25

In the foregoing and in the following examples, all temperatures are set forth uncorrected in degrees Celsius and unless otherwise indicated, all parts and percentages are by weight.

30

Examples which illustrate the invention

The content of 5-methyltetrahydrofolic acid salt which is quoted in the examples was determined by HPLC in each case and is given as % area. The water content was determined by a Karl Fischer method.

Example 1 [stabilities]

In order to determine the stabilities of the crystalline salts of 5-methyltetrahydrofolic acid, the substances were stored, together with comparison specimens, in air at 25°C and at 60% relative humidity. The content of 5-methyltetrahydrofolic acid salt remaining was measured at periodic intervals and is given by comparison with the initial value.

	Time of storage in months					
	0	3	6	12	18	88
Crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid	100%	98.6%	98.7%	99.1%	99.0%	97.8%
Amorphous calcium salt of 5-methyl-(6S)-tetrahydrofolic acid	100%			84.2%		

The crystalline salts of 5-methyltetrahydrofolic acid were still very light in colour even after an extended period of storage. In contrast thereto, the amorphous samples exhibited considerable discoloration, which occurred very rapidly.

Example 2 [X-ray powder plots]

X-ray powder plots (diffraction spectra) of these substances were recorded in order to characterise the structural properties (crystalline modifications) of the crystalline salts of 5-methyltetrahydrofolic acid.

The crystalline salts of 5-methyltetrahydrofolic acid exhibited spectra of good resolution, with sharp bands and low background effects. The spectra indicated highly crystalline constituents.

Examples of spectra are illustrated in Figure 1 (Type I), Figure 2 (Type II), Figure 3 (Type III) and Figure 4 (Type IV), and are presented in Table 1 (Type I), Table 2 (Type II), Table 3 (Type III) and Table 4 (Type IV). For comparison, a spectrum of an amorphous sample was also recorded under analogous conditions and is presented as Figure 5 (amorphous).

Selected 2 theta values for the different crystalline modifications of the crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid are listed below:

Type	Selected 2 theta values
Type I	6.5, 13.3, 16.8 and 20.1
Type II	5.3, 6.9, 18.7 and 21.1
Type III	6.8, 10.2, 15.4 and 22.5
Type IV	6.6, 15.9, 20.2 and 22.5

Example 3 [solubilities]

The solubility of the crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid is given in the following Table:

Type	Solubility at 20°C in	
	0.9% NaCl	water
Type I	1.6%	1.1%
Type II	5.8%	3.8%
Type III	1.5%	1.0%

Example 4 [amorphous calcium salt of 5-methyl-(6S)-tetrahydrofolic acid]

5 7.5 g 5-methyl-(6S)-tetrahydrofolic acid were introduced into 75 ml water at room temperature whilst passing N₂ into the batch, and the batch was adjusted to pH 12 with aqueous 30% sodium hydroxide solution. The clear solution which was thus obtained was adjusted to pH 7.5 with 37% hydrochloric acid and was treated with a solution of 7.15 g calcium chloride 6H₂O in 11.7 ml water. The white suspension which
10 was formed was stirred for 5 hours and was then filtered under suction at room temperature. The solid was washed with water and was dried under vacuum at 45°C.

5.8 g of a white, amorphous calcium salt of 5-methyl-(6S)-tetrahydrofolic acid were obtained, which had a content of 98.0% and a 6S fraction corresponding to 99.6%.

15 Even after treating this substance at 60°C in a humidity cabinet, no crystalline fractions could be determined either under a polarising microscope or by X-ray diffraction measurements.

Example 5 [crystalline calcium salt of 5-methyl-(6 R,S)-tetrahydrofolic acid]

20 70 g 5-methyl-(6R, S)-tetrahydrofolic acid were placed in a vessel in 780 ml water and the batch was adjusted to pH 7.5 with 45.2 g of 30% NaOH. The clear, slightly reddish solution was treated with a solution of 62.7 g calcium chloride 6H₂O in 140 ml

water, and the solid was filtered off and washed with a little water. The crude product which was thus obtained was suspended in water and treated at 90°C for 24 hours.

- 74.0 g of a white, crystalline calcium salt of 5-methyl-(6R,S)-tetrahydrofolic acid was
5 obtained, with a content of 99.1%.

Example 6 [crystalline calcium salt of 5-methyl-(6R)-tetrahydrofolic acid]

- 10 16.5 g 5-methyl-(6R)-tetrahydrofolic acid were placed in a vessel in 100 ml water at 92°C with 50 g calcium chloride 6H₂O. The clear, slightly yellowish suspension was stirred for 10 minutes at 91°C, and the solid was filtered off, washed with a little water and dried at 35°C under vacuum.
- 15 15.4 g of a light beige crystalline calcium salt of 5-methyl-(6R)-tetrahydrofolic acid were obtained, with a content of 97.9% and a water content of 7.8%.

Example 7 [Type I]

- 20 130 kg water were placed in a vessel and 12.8 kg 5-methyl-(6S)-tetrahydrofolic acid were introduced. The pH was adjusted to 11.6 with about 9.1 kg of 30% NaOH, and was then adjusted to 7.6 with about 1.9 kg of 37% hydrochloric acid. A suspension containing 0.3 kg carbon and 0.3 kg Cellflock was added to the clear solution. The
25 solid was filtered off and washed with 13 litres of water. The filtrate was treated with a solution containing 8.3 kg calcium chloride 2H₂O, heated to 90°C and stirred for 30 minutes. The product was filtered hot and was washed with 2 x 20 kg water. The moist crude product which was thus obtained was slurried in 115 litres of water, heated to 90°C, immediately filtered hot, washed with 2 x 20 kg water, and dried at
30 40°C under vacuum.

11.6 kg of a white, crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid (Type I) were obtained, which had a purity of 99.0% and a water content of 14.5%.

5 **Example 8 [Type I]**

1600 ml water were placed in a vessel and 194 g 5-methyl-(6S)-tetrahydrofolic acid were introduced. The pH was adjusted to 7.0 with about 80 ml of 30% NaOH. A suspension containing 20 g carbon and 20 g Cellflock in 190 ml water was added to the clear solution. The solid was filtered off and washed with water. The filtrate was treated with 950 ml of a 5.5 M calcium chloride solution, heated to 90°C and stirred for 60 minutes. The product was filtered hot, washed with water, and dried at 45°C under vacuum.

15 156.2 g of a white, crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid (Type I) were obtained, with a purity of 99.7% and a 6S fraction of 99.9%.

Example 9 [Type 1 and conversion into Type II]

20 554 g water were placed in a vessel and 53.1 g 5-methyl-(6S)-tetrahydrofolic acid were introduced. The pH was adjusted to 7.5 with 30% NaOH. 1.3 g carbon, 1.3 g Cellflock and 19.5 g water were added to the clear solution. The suspension was filtered and the solid was washed with 55 ml water. The filtrate was treated with a solution of 52.0 g calcium chloride 6H₂O in 84.6 g water, and was heated to 90°C and seeded with 100 mg of the crystalline calcium salt of 5-methyltetrahydrofolic acid. After crystallisation had occurred, the product was filtered hot at 90°C and was washed with 2 x 103 g water. The moist crude product which was thus obtained was slurried in 480 ml water, heated to 90°C, immediately filtered hot, washed as above, 30 and dried at 45°C under vacuum.

47.5 g of a white, crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid (Type I) were obtained, with a purity of 98.8% and a water content of 12.2%.

This Type I modification could be converted into the Type II modification with a water content of 5.0% by drying it at 70°C under vacuum for 30 minutes.

Example 10 [Type III]

10 15.8 g of the calcium salt of 5-methyl-(6S)-tetrahydrofolic acid were heated to 95°C in 140 ml water whilst passing N₂ through the batch. After 30 minutes at 95°C the white suspension was filtered hot under suction, and the solid was washed with water and dried at 35°C under vacuum.

15 14.0 g of a white, crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid (Type III) was obtained, with a content of 98.9% and a 6S fraction of 99.9%.

Example 11 [Type IV]

20 20.0 g of the calcium salt of 5-methyl-(6S)-tetrahydrofolic acid were heated to 100°C in 180 ml water whilst passing N₂ through the batch. After 30 minutes at 100°C the white suspension was filtered hot under suction, and the solid was washed with water and dried at 25°C under vacuum.

25 16.9 g of a white, crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid (Type IV) were obtained, with a content of 98.3% and a water content of 9.9%.

30 By drying it at 65°C under vacuum, the water content of this product could be reduced to 5.5% without a different crystalline modification being obtained in the course of this procedure.

Examples of Uses of the Crystalline Material

(a) As constituent for the production of drugs:

Beside the well known uses of 5-formyltetrahydrofolic acid previously described, reference is invited to U.S. Serial Number 09/095,572 for pharmaceutical preparations including pharmaceutically acceptable excipients, as well as dosage ranges of 5-methyltetrahydrofolic acid.

(b) As food additive:

The range of possible applications as food additive is very wide (from single preparations to multivitamin preparations and grain to breakfast foods). Please refer to e.g. WO 97/27764, SAMSF/Bailey et al., for examples (e.g. examples 1, 2, 6, 8 and 9 all contain formulas with 5-methyltetrahydrofolic acid).

(c) For the production of other tetrahydrofolic acid derivatives:

Under "other tetrahydrofolic acid derivatives", other salts of 5-methyltetrahydrofolic acid and the free acid itself are especially included.

The preceding examples can be repeated with similar success by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples. Also, the preceding specific embodiments are to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The entire disclosure of all applications, patents and publications, cited above and below, and of corresponding Swiss application 695/99, are hereby incorporated by reference.

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

Claims

- 1 Crystalline salts of 5-methyl-(6R,S)-, -(6S)- and -(6R)-tetrahydrofolic acid.
- 5 2 Crystalline salts of 5-methyl-(6S)- and -(6R)-tetrahydrofolic acid.
- 3 The crystalline calcium salt of 5-methyl-(6S)- and -(6R)-tetrahydrofolic acid.
- 4 The crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid with 2 theta
10 values of 6.5, 13.3, 16.8 and 20.1.
- 5 The crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid with 2 theta
values of 5.3, 6.9, 18.7 and 21.1.
- 15 6 The crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid with 2 theta
values of 6.8, 10.2, 15.4 and 22.5.
- 7 The crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid with 2 theta
values of 6.6, 15.9, 20.2 and 22.5.
- 20 8 A method of producing crystalline salts of 5-methyl-(6R,S)-, -(6S)- and
5-methyl-(6R)-tetrahydrofolic acid, characterised in that salts of 5-methyl-
(6R,S)-, -(6S)- or -(6R)-tetrahydrofolic acid are crystallised from a polar medium
after thermal treatment.
- 25 9 A method according to claim 8, characterised in that crystallisation is effected
after thermal treatment at a temperature above 60°C.
- 10 10 A method according to claim 8, characterised in that crystallisation is effected
30 after thermal treatment at a temperature above 85°C.

- 11 A method according to claim 8, characterised in that crystallisation is effected from a solution.
- 12 A method according to claim 8, characterised in that crystallisation is effected from a suspension.
- 13 A method according to claims 11 or 12, characterised in that crystallisation is effected from water or from a mixture of water and an organic solvent which is miscible with water.
- 14 The use of crystalline salts of 5-methyl-(6S)- or -(6R)-tetrahydrofolic acid as constituent for the production of drugs or as a food additive.
- 15 Preparations containing crystalline salts of 5-methyl-(6S)- or -(6R)-tetrahydrofolic acid.

Abstract

This invention relates to stable crystalline salts of 5-methyl-(6R,S)-, -(6S)- and -(6R)-
tetrahydrofolic acid, to methods of producing these salts and to the use thereof use
5 as a constituent for the production of drugs or as a food additive, and to preparations
containing these salts.

100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120

Total Counts







Figure 1: Crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid (Type I)

Total Counts







Total Counts




Table 1: Crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid (Type I)

```

Diffractometer : Transmission
Monochromator  : Curved Ge(111)
Wavelength     : 1.540598 Cu
Detector       : Linear PSD
Scan Mode      : Debye-Scherrer / Moving PSD / Fixed omega
2Theta scan

! Peak search parameters : Expected halfwidth : .150
!                         : Significance level : 2.5
!                         : Peak height level : 10

Peaklist [ Range 1 : 2Theta = 5.000 34.980 .020 Imax = 765 ]

!      D      2Theta  I(rel)  I(abs)  FWHM      h      k      l
13.474630    6.5544   100.0    755    .2200
8.979750     9.8420    18.5    140    .1600
6.936035    12.7526    20.3    153    .1600
6.662427    13.2786    38.3    289    .0800
6.497896    13.6164    29.4    222    .1200
6.323536    13.9935    18.8    142    .0200
6.148863    14.3933    14.0    106    .0400
5.966675    14.8352    15.5    117    .1200
5.593548    15.8309    27.5    208    .2200
5.368022    16.5006    19.7    149    .1127
5.282104    16.7709    42.5    321    .2000
4.977751    17.8044    23.6    178    .1800
4.672452    18.9782    32.7    247    .2600
4.411916    20.1102    34.8    263    .0800
4.257688    20.8467    34.2    258    .2600
3.761157    23.6360    13.3    100    .0400
3.699455    24.0361    22.3    168    .1400
3.558431    25.0037    14.8    112    .1000
3.439070    25.8864    21.0    159    .1400
3.272550    27.2283    22.1    167    .2600
3.218939    27.6907    17.0    129    .1400
3.140884    28.3931    17.2    130    .0600
3.013536    29.6198    13.9    105    .1000
2.873482    31.0991    15.1    114    .0200
2.782802    32.1395    16.6    125    .0200
2.754830    32.4748    20.2    152    .0600
2.713309    32.9858    15.4    116    .1127

```

Table 2: Crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid (Type II)

Diffractometer : Transmission
 Monochromator : Curved Ge(111)
 Wavelength : 1.540598 Cu
 Detector : Linear PSD
 Scan Mode : Debye-Scherrer / Moving PSD / Fixed omega
 2Theta scan

! Peak search parameters : Expected halfwidth : .150
 ! Significance level : 2.5
 ! Peak height level : 10

Peaklist [Range 1 : 2Theta = 5.000 34.980 .020 Imax = 526]

!	D	2Theta	I(rel)	I(abs)	FWHM	h	k	l
	12.720530	6.9434	100.0	517	.2600			
	8.508053	10.3891	29.4	152	.2400			
	6.631466	13.3409	19.6	101	.1200			
	5.883504	15.0461	71.2	368	.2200			
	5.580025	15.8696	27.8	144	.0800			
	5.010988	17.6854	42.5	220	.1400			
	4.730443	18.7434	53.6	277	.1400			
	4.215807	21.0561	35.5	184	.0400			
	3.943879	22.5263	38.8	201	.3600			
	3.581969	24.8368	24.8	128	.0200			
	3.493985	25.4726	29.6	153	.0400			
	3.309171	26.9212	22.7	117	.0200			

Table 3: Crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid (Type III)

Diffractometer : Transmission
 Monochromator : Curved Ge(111)
 Wavelength : 1.540598 Cu
 Detector : Linear PSD
 Scan Mode : Debye-Scherrer / Moving PSD / Fixed omega
 2Theta scan

! Peak search parameters : Expected halfwidth : .150
 ! Significance level : 2.5
 ! Peak height level : 10

Peaklist [Range 1 : 2Theta = 5.000 34.980 .020 Imax = 817]

!	D	2Theta	I(rel)	I(abs)	FWHM	h	k	l
	12.933490	6.8289	100.0	786	.1200			
	11.036740	8.0043	18.9	149	.0400			
	9.945525	8.8842	18.4	145	.1000			
	8.877709	9.9554	12.4	98	.0796			
	8.640580	10.2293	49.6	390	.1000			
	7.873330	11.2292	6.4	50	.1000			
	7.144004	12.3799	7.6	59	.0800			
	6.948557	12.7295	20.3	159	.1000			
	6.659956	13.2835	10.1	80	.0400			
	6.466239	13.6834	7.6	60	.0200			
	6.305060	14.0349	37.6	296	.1000			
	6.154434	14.3802	16.4	129	.0400			
	6.057193	14.6123	15.3	121	.0600			
	5.920458	14.9517	17.6	139	.1000			
	5.738533	15.4285	48.9	385	.1000			
	5.530167	16.0136	30.3	238	.1000			
	5.322477	16.6428	18.1	143	.0600			
	5.245302	16.8894	47.4	372	.0800			
	5.154604	17.1688	20.9	164	.0796			
	5.038273	17.5888	30.8	242	.1000			
	4.980502	17.7945	10.7	84	.0796			
	4.759336	18.6286	31.6	248	.1200			
	4.702846	18.8544	24.3	191	.0796			
	4.575841	19.3827	15.6	122	.0800			
	4.478961	19.8061	25.9	204	.1000			
	4.377158	20.2716	48.1	378	.1000			
	4.309006	20.5957	11.9	93	.0796			
	4.242777	20.9207	31.3	246	.0800			
	4.051441	21.9207	10.3	81	.0200			
	3.940356	22.5467	67.8	533	.1200			
	3.782452	23.5010	12.4	98	.0400			
	3.609291	24.6458	9.5	75	.0200			
	3.523157	25.2582	27.0	212	.2000			
	3.460874	25.7205	43.4	341	.0800			
	3.408545	26.1223	12.4	98	.0796			
	3.341048	26.6596	16.1	127	.2000			
	3.273575	27.2196	28.4	223	.1400			
	3.188038	27.9645	12.6	99	.0200			
	3.160110	28.2168	12.5	98	.0400			
	3.103472	28.7427	15.0	118	.0800			
	3.052658	29.2317	13.9	109	.0600			
	3.017419	29.5808	27.7	218	.1400			
	2.970195	30.0621	10.6	83	.1200			
	2.921067	30.5800	13.9	109	.0200			
	2.899222	30.8161	9.6	76	.0796			
	2.870572	31.1314	9.6	75	.0400			
	2.830661	31.5817	11.0	86	.0200			
	2.758126	32.4349	11.3	89	.0400			
	2.733265	32.7382	13.2	104	.0600			
	2.695836	33.2058	13.7	108	.0800			
	2.660160	33.6643	11.7	92	.1000			
	2.609572	34.3369	9.2	72	.0200			

Table 4: Crystalline calcium salt of 5-methyl-(6S)-tetrahydrofolic acid (Type IV)

Diffractometer : Transmission
 Monochromator : Curved Ge(111)
 Wavelength : 1.540598 Cu
 Detector : Linear PSD
 Scan Mode : Debye-Scherrer / Moving PSD / Fixed omega
 2Theta scan

! Peak search parameters : Expected halfwidth : .150
 ! Significance level : 2.5
 ! Peak height level : 10

Peaklist [Range 1 : 2Theta = 5.000 34.980 .020 Imax = 473]

!	D	2Theta	I(rel)	I(abs)	FWHM	h	k	l
	13.398610	6.5916	97.7	446	.1600			
	12.930100	6.8307	100.0	457	.0915			
	11.033220	8.0069	19.2	88	.0800			
	9.952926	8.8776	16.7	76	.1200			
	8.912272	9.9167	25.5	116	.1600			
	8.626970	10.2455	48.9	223	.0800			
	6.931997	12.7600	37.4	171	.1000			
	6.651761	13.3000	39.7	181	.1200			
	6.499623	13.6127	32.8	150	.0800			
	6.309299	14.0254	47.0	215	.1600			
	6.161306	14.3641	25.1	115	.1200			
	5.917463	14.9593	27.0	124	.1000			
	5.736254	15.4347	49.8	227	.0800			
	5.544314	15.9724	36.7	169	.1600			
	5.255854	16.8553	62.1	284	.2400			
	5.172075	17.1303	29.5	135	.0915			
	5.035719	17.5978	37.0	169	.1200			
	4.978813	17.8006	31.3	143	.0400			
	4.758441	18.6321	40.7	186	.1000			
	4.688853	18.9112	46.0	210	.0915			
	4.577465	19.3757	29.5	135	.0915			
	4.479376	19.8043	35.5	162	.1000			
	4.383704	20.2410	63.6	290	.1200			
	4.246196	20.9037	59.5	272	.1400			
	4.088125	21.7216	19.7	90	.0200			
	3.941748	22.5386	62.9	288	.1400			
	3.778991	23.5229	27.9	128	.0400			
	3.696576	24.0551	30.5	139	.1000			
	3.523769	25.2537	35.6	163	.2400			
	3.459683	25.7295	44.7	204	.0800			
	3.338511	26.6803	28.7	131	.0200			
	3.273450	27.2206	45.5	208	.1000			
	3.135320	28.4446	23.6	108	.0600			
	3.108154	28.6985	25.9	118	.0200			
	3.018687	29.5681	34.4	157	.1400			
	2.923031	30.5589	21.9	100	.0200			
	2.844431	31.4249	18.4	84	.0200			
	2.749393	32.5408	28.5	130	.1200			
	2.713739	32.9804	25.6	117	.0200			
	2.663207	33.6246	19.6	90	.0600			
	2.613490	34.2838	17.4	80	.0200			